

Algorithmic Computation of Polynomial Amoebas

D. V. Bogdanov¹, A. A. Kytmanov², and T. M. Sadykov¹

¹ Plekhanov Russian University
Stremyanny 36, Moscow, 125993, Russia
`Sadykov.TM@rea.ru`

² Siberian Federal University
Svobodny 79, Krasnoyarsk, 660041, Russia
`aakyt@gmail.com`

Abstract. We present algorithms for computation and visualization of polynomial amoebas, their contours, compactified amoebas and sections of three-dimensional amoebas by two-dimensional planes. We also provide a method and an algorithm for the computation of polynomials whose amoebas exhibit the most complicated topology among all polynomials with a fixed Newton polytope. The presented algorithms are implemented in computer algebra systems Matlab 8 and Mathematica 9.

Keywords: amoebas, Newton polytope, optimal algebraic hypersurface, the contour of an amoeba, hypergeometric functions

1 Introduction

The amoeba of a multivariate polynomial in several complex variables is the Reinhardt diagram of its zero locus in the logarithmic scale with respect to each of the variables [17]. The term "amoeba" has been coined in [5, Chapter 6] where two competing definitions of the amoeba of a polynomial have been given: the affine and the compactified versions. Both definitions are only interesting in dimension two and higher since the amoeba of a univariate polynomial is a finite set that can be explored by a variety of classical methods of localization of polynomial roots.

The geometry of the amoeba of a polynomial carries much information on the zeros of this polynomial and is closely related to the combinatorial structure of its Newton polytope (see Theorem 1). Despite losing half of the real dimensions, the image of the zero locus of a polynomial in the amoeba space reflects the relative size of some of its coefficients.

From the computational point of view, the problem of giving a complete geometric or combinatorial description of the amoeba of a polynomial is a task of formidable complexity [8, 11], despite the substantial recent progress in this direction [12–14]. The number of connected components of an amoeba complement as a function of the coefficients of the polynomial under study is still to be explored by means of the modern methods of computer algebra. In particular,

the conjecture by M.Passare on the solidness of maximally sparse polynomials (see Definition 5) remains open for a long time.

Amoebas can be computed and depicted by means of a variety of approaches and methods (see [3, 6, 9, 11, 13, 14] and the references therein). The present paper is meant to move forward the art of computing and drawing complex amoebas of algebraic hypersurfaces. A special attention paid to the geometrically most interesting and computationally most challenging case of optimal hypersurfaces (see Definition 4). We expose methods and algorithms for computation and visualization of amoebas of bivariate polynomials, their contours and compactified versions. The developed algorithms are used in higher dimensions for depicting sections of amoebas of polynomials in three variables. The main focus of the paper is on polynomials whose amoebas have the most complicated topological structure among all polynomials with a given Newton polytope (see Definition 4). We provide an algorithm for explicit construction of such polynomials.

The presented algorithms are implemented in the computer algebra systems Matlab 8 (64-bit) and Wolfram Mathematica 9 (64-bit). All examples in the paper have been computed on Intel Core i5-4440 CPU clocked at 3.10 GHz with 16 Gb RAM under MS Windows 7 Ultimate SP1.

2 Convex Polytopes, Cones and Amoebas: Definitions and Preliminaries

Let $p(x)$ be a polynomial in n complex variables:

$$p(x_1, \dots, x_n) = \sum_{\alpha \in A} c_{\alpha} x^{\alpha} = \sum_{\alpha \in A} c_{\alpha_1 \dots \alpha_n} x_1^{\alpha_1} \cdot \dots \cdot x_n^{\alpha_n},$$

where $A \subset \mathbb{Z}^n$ is a finite set.

Definition 1. *The Newton polytope $\mathcal{N}_{p(x)}$ of a polynomial $p(x)$ is the convex hull of the set A of its exponent vectors.*

Definition 2. *The recession cone of a convex set M is the set-theoretical maximal element in the family of convex cones whose shifts are contained in M .*

Definition 3. *The amoeba $\mathcal{A}_{p(x)}$ of a polynomial $p(x)$ is the image of its zero locus under the map $\text{Log} : (x_1, \dots, x_n) \mapsto (\ln |x_1|, \dots, \ln |x_n|)$.*

The connected components of the complement to the amoeba $\mathcal{A}_{p(x)}$ are convex and in bijective correspondence with the expansions of the rational function $1/p(x)$ into Laurent series centered at the origin [4]. The next statement shows how the Newton polytope $\mathcal{N}_{p(x)}$ is reflected in the geometry of the amoeba $\mathcal{A}_{p(x)}$ [4, Theorem 2.8 and Proposition 2.6].

Theorem 1. (See [4].) Let $p(x)$ be a Laurent polynomial and let $\{M\}$ denote the family of connected components of the amoeba complement ${}^c\mathcal{A}_{p(x)}$. There exists an injective function $\nu : \{M\} \rightarrow \mathbb{Z}^n \cap \mathcal{N}_{p(x)}$ such that the cone that is

dual to the polytope $\mathcal{N}_{p(x)}$ at the point $\nu(M)$ coincides with the recession cone of M . In particular, the number of connected components of ${}^c\mathcal{A}_{p(x)}$ cannot be smaller than the number of vertices of the polytope $\mathcal{N}_{p(x)}$ and cannot exceed the number of integer points in $\mathcal{N}_{p(x)}$.

Thus the amoeba of polynomial $p(x)$ in $n \geq 2$ variables is a closed connected unbounded subset of \mathbb{R}^n whose complement consists of a finite number of convex connected components. Besides, a two-dimensional amoeba has "tentacles" that go off to infinity in the directions that are orthogonal to the sides of the polygon $\mathcal{N}_{p(x)}$ (see Fig. 2, 3).

The two extreme values for the number of connected components of an amoeba complement are of particular interest.

Definition 4. (See [4, Definition 2.9].) An algebraic hypersurface $\mathcal{H} \subset (\mathbb{C}^*)^n$, $n \geq 2$, is called *optimal* if the number of connected components of the complement of its amoeba $\mathcal{A}_{\mathcal{H}}$ equals the number of integer points in the Newton polytope of the defining polynomial for \mathcal{H} . We will say that a polynomial (as well as its amoeba) is optimal if its zero locus is an optimal algebraic hypersurface.

In other words, an algebraic hypersurface is optimal if the topology of its amoeba is as complicated as it could possibly be under the condition that the Newton polytope of the defining polynomial is fixed. The other extreme case of the topologically simplest possible amoeba is defined as follows.

Definition 5. (See [10].) An algebraic hypersurface $\mathcal{H} \subset (\mathbb{C}^*)^n$, $n \geq 2$, is called *solid*, if the number of connected components of its amoeba complement ${}^c\mathcal{A}_{\mathcal{H}}$ equals the number of vertices of the Newton polytope of its defining polynomial \mathcal{H} .

Thus the solid and the optimal amoebas are the endpoints of the spectrum of amoebas of polynomials with a given Newton polytope. Of course there exist plenty of optimal (as well as solid) amoebas defined by polynomials with a given Newton polytope. In fact, both sets of amoebas regarded as subsets in the complex space of coefficients of defining polynomials, have nonempty interior.

In the bivariate case, an amoeba is solid if and only if all of the connected components of its complement are unbounded and no its tentacles are parallel. A two-dimensional optimal amoeba has, on the contrary, the maximal possible number of bounded connected components in its complement and the maximal possible number of parallel tentacles.

The functional dependency of the topological type of the amoeba $\mathcal{A}_{p(x)}$ on the coefficients of its defining polynomial $p(x)$ is complex and little understood at the present moment. A sufficient condition for the amoeba of a polynomial to be optimal is that it satisfies a "natural" system of partial differential equations of hypergeometric type [1, 2] while the support of the polynomial in question is complex enough [15]. In Section 4 we expose an algorithm for the computation of the hypergeometric polynomial with the prescribed Newton polytope.

Example 1. Let \mathcal{N} denote the convex hull of the set of lattice points $\{(0, 0), (1, 0), (0, 2), (2, 1)\}$, see Fig. 1. This polygon will appear in several examples that follow and has been chosen as one of the simplest polygons that contain an inner integer point as well as an integer point in the relative interior of its edge.




Fig. 1: The support of the polynomials in Example 1

Fig. 2 shows the amoebas of the four bivariate polynomials $p_1(x, y) = 1 + x + y + xy + y^2 + x^2y$, $p_2(x, y) = 1 + x + 3y + xy + y^2 + x^2y$, $p_3(x, y) = 1 + x + y + 4xy + y^2 + x^2y$, $p_4(x, y) = 1 + x + 3y + 4xy + y^2 + x^2y$ whose Newton polygons coincide with \mathcal{N} . The complement of the solid amoeba in Fig. 2 (a) consists of the four unbounded connected components with two-dimensional recession cones. The complement to the optimal amoeba in Fig. 2 (d) comprises six connected components: the four unbounded components with the two-dimensional recession cones, one unbounded component between the tentacles with the one-dimensional recession cone and the bounded component. The other two amoebas depicted in Fig. 2 (b) and 2 (c) exhibit five connected components in their respective complements and topologically assume an intermediate position between the solid and the optimal amoebas defined by polynomials with the Newton polygon \mathcal{N} .

The existing analytic methods [4, 12] do not in general allow one to predict the topological type of a polynomial with generic coefficients. From the computational point of view the tasks of depicting the amoebas in Fig. 2 (a) and 2 (d) are rather similar. Yet, to predict the existence of a bounded connected component of a given order [4] in an amoeba complement by means of analytic methods is in general a task of formidable complexity [11].

3 Computing Two-Dimensional Amoebas

Definition 6. We will call the "carcase" of an amoeba \mathcal{A} any subset of \mathcal{A} , such that the number of connected components of the complement to the intersection $\mathcal{A} \cap B$ for a sufficiently large ball B is as big as it could possibly be (that is, equal to the number of connected components in the complement of \mathcal{A} in \mathbb{R}^n).

We remark that the carcase of an amoeba is not uniquely defined. However, the topology of its complement in a sufficiently big ball is well-defined and is as complex as possible. When speaking of depicting an amoeba we will mean depicting its suitable carcase.

In this section we consider bivariate polynomials of the form $p(x, y) = \sum c_{ij}x^i y^j$ and provide an algorithm for computing their amoebas.

Theorem 1 yields that the geometry of the amoeba \mathcal{A}_p is closely related to the properties of the Newton polytope \mathcal{N}_p of the polynomial p . Yet, the coefficients of p also play a role and determine the size of the carcase of the amoeba in question. In what follows, the boundary of the domain where the carcase of an amoeba is depicted has been determined experimentally.

Algorithm 1 Algorithm for computing the amoeba \mathcal{A}_p of a bivariate polynomial

Require: List of the polynomial coefficients cx_list , in x by all monomials y^k , $k = 0, \dots, \deg_y p$; the boundaries of the rectangular domain in the logarithmic amoeba space a, b ; the number of values of the absolute value n_r and the angle n_φ of the complex variable.

Ensure: List of coordinates of points that belong to the amoeba carcase z_list, w_list .

```

1: procedure AMOEBA2D( $cx\_list, a, b, n_r, n_\varphi$ )
2:    $z\_list :=$  empty list
3:    $w\_list :=$  empty list
4:    $d :=$  the number of elements in  $cx\_list - 1$ 
5:    $1d := (1, \dots, 1) \in \mathbb{R}^d$   $\triangleright$  the vector with  $d$  units
6:    $h_r := (\exp(b) - \exp(a)) / (n_r - 1)$ 
7:    $h_\varphi := 2\pi / (n_\varphi - 1)$ 
8:   for  $r = \exp(a) : \exp(b) : h_r$  do
9:     for  $\varphi = 0 : 2\pi : h_\varphi$  do
10:       $x := r * \exp(\sqrt{-1} * \varphi)$ 
11:       $y := \text{roots}(cx\_list)$   $\triangleright$  the vector of zeros of the polynomial with the
        coefficients  $cx\_list$ 
12:      Add  $\text{Log}(|x| * 1d)$  to  $z\_list$ 
13:      Add  $\text{Log}(|y|)$  to  $w\_list$ 
14:    end for
15:  end for
16:  return  $\{z\_list, w\_list\}$ 
17: end procedure

```

The next straightforward algorithm which computes the amoeba of a bivariate polynomial has been used by numerous authors in various forms [3, 6, 9]. We include it for the sake of completeness and future reference. We refer to [8] for a detailed discussion of computational complexity of polynomial root finding algorithms.

To obtain a picture of good quality the steps of the Algorithm 1 are repeated with the variables x and y interchanged. The points with the computed coordinates are depicted in the same figure.

Example 2. For the polynomial $p_4(x, y)$ in Example 1 the lists of polynomial coefficients in the variables x and y are as follows: $cx_list = \{1, 3 + 4x + x^2, 1 + x\}$ and $cy_list = \{y, 1 + 4y, 1 + 3y + y^2\}$. The (carcase of the) amoeba is depicted in the rectangle $[-5, 5] \times [-5, 5]$. The number of values of the absolute value of a variable is $n_r = 2000$ while the number of values of its argument is $n_\varphi = 180$. Fig. 2 features the amoebas of the polynomials in Example 1 computed by means of Algorithm 1.

The topologically more involved amoeba of the polynomial in Example 4 has been computed in a similar way.




Fig. 2: The amoebas of the polynomials $p_1(x, y)$, $p_2(x, y)$, $p_3(x, y)$ and $p_4(x, y)$

4 Generating Optimal Polynomials

In this section we employ the notion of a hypergeometric polynomial for the purpose of constructive generation of optimal amoebas. We will need the following auxiliary definition.

Definition 7. A formal Laurent series

$$\sum_{s \in \mathbb{Z}^n} \varphi(s) x^s \quad (1)$$

is called *hypergeometric* if for any $j = 1, \dots, n$ the quotient $\varphi(s + e_j)/\varphi(s)$ is a rational function in $s = (s_1, \dots, s_n)$. Throughout the paper we denote this rational function by $P_j(s)/Q_j(s + e_j)$. Here $\{e_j\}_{j=1}^n$ is the standard basis of the lattice \mathbb{Z}^n . By the *support* of this series we mean the subset of \mathbb{Z}^n on which $\varphi(s) \neq 0$.

A *hypergeometric function* is a (multi-valued) analytic function obtained by means of analytic continuation of a hypergeometric series with a nonempty domain of convergence along all possible paths [5, 10].

Theorem 2. (Ore, Sato, cf [16]) *The coefficients of a hypergeometric series are given by the formula*

$$\varphi(s) = t^s U(s) \prod_{i=1}^m \Gamma(\langle \mathbf{A}_i, s \rangle + c_i), \quad (2)$$

where $t^s = t_1^{s_1} \dots t_n^{s_n}$, $t_i, c_i \in \mathbb{C}$, $\mathbf{A}_i = (A_{i,1}, \dots, A_{i,n}) \in \mathbb{Z}^n$, $i = 1, \dots, m$ and $U(s)$ is the product of a certain rational function and a periodic function $\phi(s)$ such that $\phi(s + e_j) = \phi(s)$ for every $j = 1, \dots, n$.

Given the above data $(t_i, c_i, \mathbf{A}_i, U(s))$ that determines the coefficient of a multivariate hypergeometric Laurent series, it is straightforward to compute the rational functions $P_i(s)/Q_i(s + e_i)$ using the Γ -function identity $\Gamma(z+1) = z\Gamma(z)$ (see e.g. [16]).

Algorithm 2 Generation of a bivariate hypergeometric polynomial and its defining system of equations

Require: List of vertices N_list of a convex integer polygon P .

Ensure: List of coefficients c_list of the hypergeometric polynomial $p(x, y)$ whose Newton polygon is P .

```

1: procedure HYPERPOLY2D( $N\_list$ )
2:    $B$  = list of outer normals to the sides of  $P$ 
3:    $\varphi(s, t) = 1 / \prod_{j=1}^q \Gamma(1 - \langle B_j, (s, t) \rangle - c_j)$ 
4:    $c\_list$  = list of the coefficients of the hypergeometric polynomial
5:    $R_1 = \text{FunctionExpand}[\varphi(s+1, t) / \varphi(s, t)]$ 
6:    $R_2 = \text{FunctionExpand}[\varphi(s, t+1) / \varphi(s, t)]$ 
7:    $P_1 = \text{Numerator}[R_1]$ 
8:    $P_2 = \text{Numerator}[R_2]$ 
9:    $Q_1 = \text{Denominator}[R_1]$ 
10:   $Q_2 = \text{Denominator}[R_2]$ 
11:   $\theta_x = xp'_x$ 
12:   $\theta_y = yp'_y$ 
13:   $p$  = the polynomial defined by  $c\_list$ 
14:  if  $xP_1(\theta)p = Q_1(\theta)p$  and  $yP_2(\theta)p = Q_2(\theta)p$  then
15:    return  $\{c\_list\}$ 
16:  end if
17: end procedure

```

Definition 8. A (formal) Laurent series $\sum_{s \in \mathbb{Z}^n} \varphi(s) x^s$ whose coefficient satisfies the relations $\varphi(s + e_j) / \varphi(s) = P_j(s) / Q_j(s + e_j)$ is a (formal) solution to the following system of partial differential equations of hypergeometric type:

$$x_j P_j(\theta) f(x) = Q_j(\theta) f(x), \quad j = 1, \dots, n. \quad (3)$$

Here $\theta = (\theta_1, \dots, \theta_n)$, $\theta_j = x_j \frac{\partial}{\partial x_j}$.

The system (3) will be referred to as *the Horn hypergeometric system defined by the Ore-Sato coefficient* $\varphi(s)$ (see [2]) and denoted by $\text{Horn}(\varphi)$.

For a convex integer polytope \mathcal{N} , we construct the list B of the outer normals to the faces of \mathcal{N} . We denote the length of this list by q . We assume the elements of B to be normalized so that the coordinates of each outer normal are integer and relatively prime. Define the Ore-Sato coefficient

$$\varphi(s) = \frac{1}{\prod_{j=1}^q \Gamma(1 - \langle B_j, s \rangle - c_j)}. \quad (4)$$

The hypergeometric system (3) defined by the Ore-Sato coefficient $\varphi(s)$ admits a polynomial solution with several interesting properties. Under the additional assumption that it cannot be factored in the ring of Puiseux polynomials $\mathbb{C}[x_1^{1/d}, \dots, x_n^{1/d}]$ for any $d \in \mathbb{N}$, this polynomial turns out to have an optimal

amoeba. We will employ Algorithm 2 to generate optimal polynomials of hypergeometric type.

Example 3. The outer normals (normalized as explained above) to the sides of the polygon shown in Fig. 1 are as follows: $(0, -1)$, $(-1, 0)$, $(1, 2)$, $(1, -1)$. Using (4) we obtain

$$\varphi(s, t) = (\Gamma(s+1) \Gamma(t+1) \Gamma(-s-2t+5) \Gamma(-s+t+2))^{-1}.$$

By Definition 8 the above Ore-Sato coefficient $\varphi(s, t)$ gives rise to the polynomials

$$\begin{aligned} P_1(s, t) &= (-s-2t+4)(-s+t+1), & Q_1(s, t) &= s+2, \\ P_2(s, t) &= (s+2t-4)(s+2t-3), & Q_2(s, t) &= (t+2)(-s+t+3). \end{aligned}$$

The corresponding hypergeometric system is defined by the linear partial differential operators

$$\begin{cases} x(-\theta_x - 2\theta_y + 4)(-\theta_x + \theta_y + 1) - (\theta_x + 2), \\ y(\theta_x + 2\theta_y - 4)(\theta_x + 2\theta_y - 3) - (\theta_y + 2)(-\theta_x + \theta_y + 3). \end{cases} \quad (5)$$

It is straightforward to check that the hypergeometric polynomial $p(x, y) = 1 + 4x + 6y + 24xy + 12x^2y + 2y^2$ (whose coefficients are found in accordance with Step 3 of Algorithm 2 and normalized to be relatively prime integers) belongs to the kernels of the operators (5).

Example 4. Using Algorithms 1 and 2 we compute the coefficients of the hypergeometric polynomial supported in the polygon $\mathcal{N}_{p_h(x,y)}$ shown in Fig. 3 (a): $p_h(x, y) = 2x^2 + 20xy + 72x^2y + 20x^3y + 5y^2 + 160xy^2 + 450x^2y^2 + 160x^3y^2 + 5x^4y^2 + 12y^3 + 300xy^3 + 800x^2y^3 + 300x^3y^3 + 12x^4y^3 + 5y^4 + 160xy^4 + 450x^2y^4 + 160x^3y^4 + 5x^4y^4 + 20xy^5 + 72x^2y^5 + 20x^3y^5 + 2x^2y^6$. The amoeba $\mathcal{A}_{p_h(x,y)}$ of this polynomial is depicted in Fig.3 (a) (right).

A more involved hypergeometric polynomial is given by $p_{h2}(x, y) = -456456x^3 + 488864376x^2y - 28756728x^3y + 25420947552x^2y^2 - 244432188x^3y^2 + 3003x^4y^2 - 119841609888xy^3 + 127104737760x^2y^3 - 465585120x^3y^3 + 6006x^4y^3 + 1396755360y^4 - 508418951040xy^4 + 139815211536x^2y^4 - 232792560x^3y^4 + 1729x^4y^4 + 4190266080y^5 - 355893265728xy^5 + 41611670100x^2y^5 - 29628144x^3y^5 + 57x^4y^5 + 698377680y^6 - 58663725120xy^6 + 3328933608x^2y^6 - 705432x^3y^6 - 2327925600xy^7 + 55023696x^2y^7 - 16930368xy^8$. Its defining Ore-Sato coefficient equals $\varphi(s, t) = \Gamma(s+t-4)\Gamma(-4s+t-16)\Gamma(-3s-2t-5)\Gamma(3s-t-3)\Gamma(2s+t-5)$. The amoeba of this polynomial is shown in Fig. 3 (b).

5 Computing Contours of Amoebas of Bivariate Polynomials

The boundary of an amoeba admits a rich analytic structure that is revealed in the following definition.




Fig. 3: The Newton polytope and the amoeba of the hypergeometric polynomial $p_h(x, y)$ and polynomial $p_{h2}(x, y)$

Definition 9. The *contour* of the amoeba $\mathcal{A}_{p(x)}$ is the set $\mathcal{C}_{p(x)}$ of critical points of the logarithmic map Log restricted to the zero locus of the polynomial $p(x)$, that is, the map $\text{Log} : \{p(x) = 0\} \rightarrow \mathbb{R}^n$.

The structure of the contour of an amoeba can be described in terms of the logarithmic Gauss map which is defined as follows.

Definition 10. Denote by $\text{Gr}(n, k)$ the Grassmannian of k -dimensional subspaces in \mathbb{C}^n . The *logarithmic Gauss map* [7] is defined to be the map $\gamma : \mathcal{H} \rightarrow \text{Gr}(n, k)$ that maps a regular point $x \in \text{reg } \mathcal{H}$ into the normal subspace $\gamma(x)$ to the image $\log \mathcal{H}$.

The boundary of an amoeba $\partial \mathcal{A}_{p(x)}$ is necessarily a subset of the contour $\mathcal{C}_{p(x)}$ but is in general different from it.

Knowing the structure of the contour of an amoeba is important for describing the topological structure of the amoeba complement. Experiments show that a cusp of the contour inside the "body" of the corresponding amoeba is a counterpart of a missing connected component in its complement.

One of the ways to draw the contour of an amoeba is to depict the solutions to the system of algebraic equations

$$\begin{cases} p(x, y) = 0, \\ x \frac{\partial p(x, y)}{\partial x} - uy \frac{\partial p(x, y)}{\partial y} = 0. \end{cases} \quad (6)$$

Here $u \in \mathbb{R} \cup \{\infty\}$ is a real parameter that encodes the slope of the normal line to the contour of the amoeba.

Eliminating the variables x and y out of the system (6) we obtain polynomials $s(y, u)$ and $t(x, u)$ that can be used to depict the contour of the amoeba by means of the following algorithm.

Remark 1. The lists of coordinates that are obtained at each iteration of the cycle in the Algorithm 3 might contain additional elements that do not correspond to the amoeba contour. They are sorted out by checking against the system of equations (6). All visualization algorithms are linear in the lattice parameters h_r, h_φ, h_u .

Algorithm 3 Computing the contour \mathcal{C}_p of the amoeba of a bivariate polynomial

Require: List of polynomial coefficients cx_list cy_list depending on u , by all monomials x^k , $k = 0, \dots, \deg_x t$, y^m , $m = 0, \dots, \deg_y s$; initial value u_1 , final value u_n and the step h_u .

Ensure: List of coordinates of points z_list , w_list that belong to the contour of the amoeba.

```

1: procedure CONTOUR2D( $cx\_list$ ,  $cy\_list$ ,  $u_1$ ,  $u_n$ ,  $h_u$ )
2:    $z\_list :=$  the empty list
3:    $w\_list :=$  the empty list
4:    $dx :=$  the number of elements in  $cx\_list - 1$ 
5:    $dy :=$  the number of elements in  $cy\_list - 1$ 
6:   for  $u = u_1 : u_n : h_u$  do
7:      $x := \text{roots}(cx\_list)$   $\triangleright$  the vector of zeros of the polynomial with the
       coefficients  $cx\_list$ 
8:      $y := \text{roots}(cy\_list)$   $\triangleright$  the vector of zeros of the polynomial with the
       coefficients  $cy\_list$ 
9:     Add  $\text{Log}(|x|)$  to  $z\_list$ 
10:    Add  $\text{Log}(|y|)$  to  $w\_list$ 
11:   end for
12:   return  $\{z\_list, w\_list\}$ 
13: end procedure

```

Example 5. The system of equations (6) associated with the first polynomial in Example 1 has the form

$$\begin{cases} 1 + x + y + xy + y^2 + x^2y = 0, \\ x - uy + xy - uxy + 2x^2y - ux^2y - 2uy^2 = 0. \end{cases}$$

Eliminating variables x and y yields the equations $u - 2uy + u^2y - 4uy^2 + 2u^2y^2 - 2uy^3 - u^2y^3 + 3uy^4 + 4uy^5 + u^2y^5 = 1 - y - 4y^2 - 9y^3 - 7y^4 - 4y^5$ and $-3u^2 - 3ux - 5u^2x - 2ux^2 + u^2x^2 + 3ux^3 + 5u^2x^3 + 2ux^4 + 3u^2x^4 + ux^5 + u^2x^5 = x^2 + 2x^3 + 5x^4 + 2x^5$, respectively. The lists of the coefficients that depend on u are $cx_list = \{-1 + u, 1 - 2u + u^2, 4 - 4u + 2u^2, 9 - 2u - u^2, 7 + 3u, 4 + 4u + u^2\}$ and $cy_list = \{-3u^2, -3u - 5u^2, -1 - 2u + u^2, -2 + 3u + 5u^2, -5 + 2u + 3u^2, -2 + u + u^2\}$. The parameter u assumes values in the interval $[-120, 120]$ with the step $h_u = 0,001$. The obtained contours of the amoebas of the polynomials in Example 1 have nontrivial structure and are shown in Fig. 4.

6 Computing Two-Dimensional Compactified Amoebas

Definition 11. The compactified amoeba $\overline{\mathcal{A}}_{p(x)}$ of a polynomial $p(x)$ is defined to be the image of its zero locus under the mapping

$$M(x) = \frac{\sum_{\alpha \in A} \alpha \cdot |x^\alpha|}{\sum_{\alpha \in A} |x^\alpha|} = \frac{\sum_{(\alpha_1, \dots, \alpha_n) \in A} (\alpha_1, \dots, \alpha_n) \cdot |x^{\alpha_1} \cdot \dots \cdot x^{\alpha_n}|}{\sum_{(\alpha_1, \dots, \alpha_n) \in A} |x^{\alpha_1} \cdot \dots \cdot x^{\alpha_n}|}. \quad (7)$$




Fig. 4: Contours of the amoebas of the polynomials $p_1(x, y)$, $p_2(x, y)$, $p_3(x, y)$ and $p_4(x, y)$

Numeric computation of two-dimensional compactified amoebas is similar to that of their affine counterparts (see Algorithm 1). The main computational issue is dealing with the moment map (7) instead of the logarithmic map into the affine space.

Example 6. Applying (7) to the polynomials in Example 1 yields the mapping

$$(x, y) \mapsto \mu(x, y) = \frac{(1 \cdot (0, 0) + |x| \cdot (1, 0) + |y| \cdot (0, 1) + |xy| \cdot (1, 1) + |y^2| \cdot (0, 2) + |x^2y| \cdot (2, 1))}{(|1| + |x| + |y| + |xy| + |y^2| + |x^2y|)} = \frac{(|x| + |xy| + 2|x^2y|, |y| + |xy| + 2|y^2| + |x^2y|)}{(|1| + |x| + |y| + |xy| + |y^2| + |x^2y|)}.$$

The corresponding compactified amoebas inside the Newton polygon of their defining polynomials are shown in Fig. 5.




Fig. 5: Compactified amoebas of the polynomials $p_1(x, y)$, $p_2(x, y)$, $p_3(x, y)$ and $p_4(x, y)$ inside their Newton polygon

7 Multivariate Outlook

Depicting three-dimensional amoebas represents a substantial computational challenge due to the complex geometry of their shape. We will now treat the

problem of computing sections of amoebas of polynomials in three variables by two-dimensional hyperplanes.

Consider a polynomial in three complex variables $p(x, y, z) = \sum c_{ijk} x^i y^j z^k$. To compute the section of its amoeba by the plane $|z| = \text{const}$ we fix the absolute value of z and modify Algorithm 1 by adding a cycle with respect to the values of the argument $\varphi_z \in [0; 2\pi]$.

Algorithm 4 Computing the section of the amoeba \mathcal{A}_p of a polynomial in three variables by a two-dimensional plane

Require: List of polynomial coefficients cxz_list , depending on x and z by all monomials y^k , $k = 0, \dots, \deg_y p$; bounds a, b for the rectangular domain in the amoeba space where the amoeba section is depicted; the numbers n_r and n_φ of values of the absolute value and the argument of the variable $z \in \mathbb{C}$, respectively.

Ensure: List of points that belong to the amoeba section z_list, w_list .

```

1: procedure AMOEBA3D( $cxz\_list, a, b, n_r, n_\varphi, z$ )
2:    $z\_list :=$  empty list
3:    $w\_list :=$  empty list
4:    $d :=$  the number of elements in  $cxz\_list - 1$ 
5:    $1d := (1, \dots, 1) \in \mathbb{R}^d$  ▷ the vector of  $d$  units
6:    $h_r := (\exp(b) - \exp(a)) / (n_r - 1)$ 
7:    $h_\varphi := 2\pi / (n_\varphi - 1)$ 
8:   for  $r = \exp(a) : \exp(b) : h_r$  do
9:     for  $\varphi = 0 : 2\pi : h_\varphi$  do
10:       $x := r * \exp(\sqrt{-1} * \varphi)$ 
11:       $y := \text{roots}(cxz\_list)$  ▷ the vector of zeros of the polynomial with the
        coefficients  $cx\_list$ 
12:      Add  $\text{Log}(|x| * 1d)$  to  $z\_list$ 
13:      Add  $\text{Log}(|y|)$  to  $w\_list$ 
14:     end for
15:   end for
16:   return  $\{z\_list, w\_list\}$ 
17: end procedure

```

Example 7. Using Algorithm 4 we compute a section of the amoeba of the polynomial $p(x, y, z) = 1 + 3y + y^2 + 6xy + x^2y + xyz + xyz^2$. This polynomial is one of the simplest polynomials with tetrahedral Newton polytopes that contain integer points in the interior as well as in the relative interior of faces of all positive dimensions (see Fig. 7). The section of the amoeba $\mathcal{A}_{p(x,y,z)}$ by the plane $\log |z| = 5$ is shown in Fig. 7 (right).

Example 8. We now consider a computationally more challenging optimal hypergeometric polynomial in three variables supported in a regular integer octahedron. Due to symmetry it suffices to consider its part that belongs to the positive orthant and is shown in Fig. 6 (left). We use a three-dimensional version of Algorithm 2 to compute the corresponding (uniquely determined up to scaling)

hypergeometric polynomial: $p_h(x, y, z) = x^2y^2 + 36x^2yz + 36xy^2z + 256x^2y^2z + 36x^3y^2z + 36x^2y^3z + x^2z^2 + 36xyz^2 + 256x^2yz^2 + 36x^3yz^2 + y^2z^2 + 256xy^2z^2 + 1296x^2y^2z^2 + 256x^3y^2z^2 + x^4y^2z^2 + 36xy^3z^2 + 256x^2y^3z^2 + 36x^3y^3z^2 + x^2y^4z^2 + 36x^2yz^3 + 36xy^2z^3 + 256x^2y^2z^3 + 36x^3y^2z^3 + 36x^2y^3z^3 + x^2y^2z^4$. Fig. 6 (right) shows the intersection of the amoeba $\mathcal{A}_{p_h(x,y,z)}$ with the three coordinate hyperplanes as well as with the hyperplanes $|x| = 3$, $|x| = 8$, $|y| = 6$, $|z| = 4$ in the logarithmic amoeba space. The figure space is orthogonal to the vector $(1, 1, 1)$.




Fig. 6: The part of the support of $p_h(x, y, z)$ that belongs to the positive orthant and the carcase of the corresponding part of the amoeba




Fig. 7: The Newton polytope $\mathcal{N}_{p(x,y,z)}$ and the section of the amoeba $\mathcal{A}_{p(x,y,z)}$ by the plane $\log |z| = 5$

We denote by (s, t, u) the coordinates in the amoeba space of Fig. 6 (right). The bold segments join the sections of the amoeba by the planes $t = 0$, $t = \log 6$, $u = 0$, $u = \log 4$, and the section of the amoeba by the plane $s = \log 3$. The bold curve is the "top of the box," i.e., the intersection of the amoeba with the plane $u = \log 4$.

Acknowledgements. This research was supported by the state order of the Ministry of Education and Science of the Russian Federation for Siberian Federal University (task 1.1462.2014/K), by grant of the Government of the Russian Federation for investigations under the guidance of the leading scientists of the Siberian Federal University (contract No. 14.Y26.31.0006) and by the Russian Foundation for Basic Research, project 15-31-20008-mol.a.ved.

References

1. Dickenstein, A., Sadykov, T. M.: Algebraicity of solutions to the Mellin system and its monodromy. *Doklady Mathematics* 75 (1), 80–82 (2007)
2. Dickenstein, A., Sadykov, T. M.: Bases in the solution space of the Mellin system. *Sbornik Mathematics* 198 (9-10), 1277–1298 (2007)
3. Forsberg, M.: Amoebas and Laurent Series. Doctoral Thesis presented at Royal Institute of Technology, Stockholm, Sweden. Bromma Tryck AB, ISBN 91-7170-259-8 (1998)
4. Forsberg, M., Passare, M., Tsikh, A. K.: Laurent determinants and arrangements of hyperplane amoebas. *Adv. Math.* 151, 45–70 (2000)
5. Gelfand, I. M., Kapranov, M. M., Zelevinsky, A. V.: Discriminants, resultants, and multidimensional determinants. *Mathematics: Theory & Applications*. Birkhäuser Boston, Inc., Boston, MA (1994)
6. Johansson, P.: On the Topology of the Coamoeba. Doctoral Thesis presented at Stockholm University, Sweden. US AB, ISBN 978-91-7447-933-1 (2014)
7. Kapranov, M. M.: A characterization of A-discriminantal hypersurfaces in terms of the logarithmic Gauss map. *Math. Ann.* 290, 277–285 (1991)
8. Kim, M.-H.: Computational complexity of the Euler type algorithms for the roots of complex polynomials. Thesis, City University of New York, New York (1985)
9. Nilsson, L.: Amoebas, Discriminants and Hypergeometric Functions. Doctoral Thesis presented at Stockholm University, Sweden. US AB, ISBN 978-91-7155-889-3 (2009)
10. Passare, M., Sadykov, T. M., Tsikh, A. K.: Nonconfluent hypergeometric functions in several variables and their singularities. *Compos. Math.* 141 (3), 787–810 (2005)
11. Purbhoo, K.: A Nullstellensatz for amoebas. *Duke Math. J.* 141 (3), 407–445 (2008)
12. Rullgård, H.: Stratification des espaces de polynômes de Laurent et la structure de leurs amibes (French). *Comptes Rendus de l'Academie des Sciences - Series I: Mathematics* 331 (5), 355–358 (2000)
13. Theobald, T.: Computing Amoebas. *Experiment. Math.* 11 (4), 513–526 (2002)
14. Theobald, T., de Wolff, T.: Amoebas of genus at most one. *Adv. Math.* 239, 190–213 (2013)
15. Sadykov, T. M., Tsikh, A. K.: Hypergeometric and Algebraic Functions in Several Variables (Russian). Nauka (2014)
16. Sadykov, T. M.: On a multidimensional system of hypergeometric differential equations. *Siberian Mathematical Journal* 39 (5), 986–997 (1998)
17. Viro, T.: What is an amoeba? *Notices of the AMS* 49 (8), 916–917 (2002)